

Easily Perform Prothrombin Time (PT) / INR

Plastinex® (Thromboplastin) is for use in performing the one stage Prothrombin Time (PT) and Coagulation Factor Assays which are based on a modified Prothrombin Time.



Net Contents: 20 x 3.0 mL

- Preparation of Lyophilized Rabbit Brain Thromboplastin, Calcium Salt, Buffers and Stabilizers
- Perform One Stage Prothrombin Time (PT)
- Evaluate the Extrinsic Pathways of Factor I (Fibrinogen), V, VII, & X
- Monitor Oral Coagulation Therapy

Includes the International Sensitivity Index (ISI) of the Reagent

The prothrombin time is the method of choice for monitoring oral anti-coagulation therapy. It is also a fundamental screening test for a deficiency or abnormality of extrinsic coagulation factor VII, and the factors common to both the intrinsic and extrinsic hemostatic pathways: fibrinogen, I, V, and X. When used in conjunction with deficient substrate plasma, the PT provides the basis for the determination of the percent (%) activity of specific coagulation factors.



155 Gibraltar Road
Horsham, PA 19044 USA

REF
101158



An ISO 13485 Registered Company
www.biodatacorp.com



mdi Europa GmbH
Langenhagener Str. 71
D-30855 Langenhagen
GERMANY



Alpha Laboratories
40 Parham Drive Eastleigh
SO50 4NU Hampshire
UNITED KINGDOM

